

UNSOLVABLE ONE-DIMENSIONAL LIFTING PROBLEMS FOR CONGRUENCE LATTICES OF LATTICES

JIŘÍ TŮMA AND FRIEDRICH WEHRUNG

ABSTRACT. Let S be a distributive $\{\vee, 0\}$ -semilattice. In a previous paper, the second author proved the following result:

Suppose that S is a lattice. Let K be a lattice, let $\varphi: \text{Con}_c K \rightarrow S$ be a $\{\vee, 0\}$ -homomorphism. Then φ is, up to isomorphism, of the form $\text{Con}_c f$, for a lattice L and a lattice homomorphism $f: K \rightarrow L$.

In the statement above, $\text{Con}_c K$ denotes as usual the $\{\vee, 0\}$ -semilattice of all finitely generated congruences of K .

We prove here that this statement characterizes S being a lattice.

INTRODUCTION

The Congruence Lattice Problem (CLP in short) asks whether for any distributive $\{\vee, 0\}$ -semilattice S , there exists a lattice L such that $\text{Con}_c L \cong S$. While this problem is still unsolved, many related problems have been solved. Among these, we mention the following, due to G. Grätzer and E.T. Schmidt, see [4, 5], and also [6] for a survey about this and related problems.

Theorem 1. *Let S be a finite distributive $\{\vee, 0\}$ -semilattice, let K be a finite lattice, let $\varphi: \text{Con}_c K \rightarrow S$ be a $\{\vee, 0\}$ -homomorphism. Then there are a finite lattice L , a lattice homomorphism $f: K \rightarrow L$, and an isomorphism $\alpha: \text{Con}_c L \rightarrow S$ such that $\alpha \circ \text{Con}_c f = \varphi$.*

In the statement of Theorem 1, $\text{Con}_c f$ denotes the map from $\text{Con}_c K$ to $\text{Con}_c L$ that with any congruence α of K associates the congruence of L generated by all the pairs $\langle f(x), f(y) \rangle$ where $\langle x, y \rangle \in \alpha$.

In [10], the second author proves that provided that S is a lattice, all finiteness assumptions in Theorem 1 can be dropped, that is:

Theorem 2. *Let S be a distributive lattice with zero, let K be a lattice, let $\varphi: \text{Con}_c K \rightarrow S$ be a $\{\vee, 0\}$ -homomorphism. Then φ can be “lifted”, that is, there are a lattice L , a lattice homomorphism $f: K \rightarrow L$, and an isomorphism $\alpha: \text{Con}_c L \rightarrow S$ such that $\alpha \circ \text{Con}_c f = \varphi$.*

In the result of Theorem 2, instead of lifting a distributive $\{\vee, 0\}$ -semilattice S (with respect to the Con_c functor), we lift a $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c K \rightarrow$

Date: February 1, 2008.

1991 Mathematics Subject Classification. 06B10, 06E05.

Key words and phrases. Lattice, congruence, amalgamation.

This work was completed while the first author was visiting the University of Caen. It was partly financed by the institutional grant CEZ:J13/98:113200007a and by GAUK no. 162/1999. The visit was financed by a BARRANDE program.

S . For this reason, we shall call such a statement “one-dimensional Congruence Lattice Problem”, in short 1-CLP. With this terminology, the usual CLP would have to be called 0-CLP. By replacing K by a truncated n -dimensional cube (diagram) of lattices, we can define the n -CLP, for any positive integer n . It turns out that this problem is interesting only for $n \in \{0, 1, 2\}$. Indeed, it follows from [8] that the 3-CLP holds only for trivial S —but much more is proved in [8], while the result about 3-CLP follows from a trivial (and unpublished) example of the second author. The 2-CLP is another matter (far less trivial than 3-CLP but still far easier than 1-CLP), which will be considered elsewhere.

Our main result (see Theorem A) states that for a given distributive $\{\vee, 0\}$ -semilattice S , Theorem 2 *characterizes* S being a lattice. This solves also a problem formulated by H. Dobbertin in the (yet unpublished) monograph [2], see Corollary 1.4. In fact, our approach is inspired by Dobbertin’s solution for the particular case of his own problem where S is *primely generated*, see Theorem 15 in [1]. It gives, for a distributive $\{\vee, 0\}$ -semilattice S that is not a lattice, the construction of a Boolean algebra B of size at most $2^{|S|}$ and a $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c B \rightarrow S$ that cannot be “lifted” as in Theorem 2.

Even in the particular case where $S = D$, the simplest distributive $\{\vee, 0\}$ -semilattice that is not a lattice, see Section 2, it has been an open problem, stated at the end of Section 1 in [1], whether the size of B can be reduced from $2^{\aleph_0}$ to $\aleph_1$ (without the Continuum Hypothesis). We solve this affirmatively in Theorem B. This also gives us that *there are a Boolean algebra B of size $\aleph_1$ and a $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c B \rightarrow D$ that cannot be lifted*, see Corollary 2.4.

We use standard notation and terminology. For a partially ordered set $\langle P, \leq \rangle$ and for $a \in P$, we put

$$(a] = \{x \in P \mid x \leq a\}.$$

We denote by ω the set of all natural numbers, and by ω_1 the first uncountable ordinal.

1. CHARACTERIZATION OF DISTRIBUTIVE $\{\vee, 0\}$ -SEMILATTICES WITH 1-CLP

The main lemma of this section is the following.

Lemma 1.1. *Let S be a distributive $\{\vee, 0\}$ -semilattice, let $\mathbf{a}_0, \mathbf{a}_1 \in S$ be such that the set $Q = (\mathbf{a}_0] \cap (\mathbf{a}_1]$ has no largest element.*

There are a Boolean algebra B and a $\{\vee, 0\}$ -homomorphism $\mu: B \rightarrow S$ such that the following holds:

- (a) $\mu(1) = \mathbf{a}_0 \vee \mathbf{a}_1$;
- (b) *there are no maps $\mu_0, \mu_1: B \rightarrow S$ that satisfy the following properties:*
 - (i) $\mu(x) = \mu_0(x) \vee \mu_1(x)$, for all $x \in B$,
 - (ii) μ_0 and μ_1 are order-preserving,
 - (iii) $\mu_\ell(1) \leq \mathbf{a}_\ell$, for all $\ell < 2$.

Proof. Let κ be the minimum size of a cofinal subset of Q , and pick a cofinal subset $\{\mathbf{x}_\xi \mid \xi < \kappa\}$ of Q . So κ is an infinite cardinal. We define recursively a map $f: \kappa \rightarrow \kappa$ by the rule

$$f(\alpha) = \min\{\xi < \kappa \mid \mathbf{x}_\xi \notin \text{Id}\{\mathbf{x}_{f(\beta)} \mid \beta < \alpha\}\} \quad (1.1)$$

for all $\alpha < \kappa$, where $\text{Id } X$ denotes the ideal of S generated by a subset X of S . Let $\beta < \alpha$. Then, by (1.1), $\mathbf{x}_{f(\alpha)} \notin \text{Id}\{\mathbf{x}_{f(\gamma)} \mid \gamma < \alpha\}$, so $f(\alpha) \neq f(\beta)$. Moreover,

$\mathbf{x}_{f(\alpha)} \notin \text{Id}\{\mathbf{x}_{f(\gamma)} \mid \gamma < \beta\}$, so $f(\beta) \leq f(\alpha)$, whence $f(\beta) < f(\alpha)$. So f is strictly increasing.

For $\alpha < \kappa$, we put $\mathbf{q}_\alpha = \mathbf{x}_{f(\alpha)}$ and $Q_\alpha = \text{Id}\{\mathbf{q}_\beta \mid \beta < \alpha\}$. By (1.1), $\mathbf{q}_\alpha \notin Q_\alpha$ for all $\alpha < \kappa$. Furthermore, all the sets Q_α are ideals of Q and $Q_\alpha \subset Q_\beta$ whenever $\alpha < \beta$. Finally, for $\alpha < \beta$, the relation $\mathbf{q}_\alpha \in Q_\beta$ holds. (Otherwise $\mathbf{x}_{f(\alpha)} \notin Q_\beta = \text{Id}\{\mathbf{x}_{f(\gamma)} \mid \gamma < \beta\}$, thus, by (1.1), $f(\beta) \leq f(\alpha)$, a contradiction since f is strictly increasing.) Hence $\bigcup_{\alpha < \kappa} Q_\alpha = Q$.

For $\mathbf{x} \in Q$, we denote by $\|\mathbf{x}\|$ the least $\alpha < \kappa$ such that $\mathbf{x} \in Q_\alpha$. Observe that the following obvious properties hold:

$$\|\mathbf{q}_\alpha\| = \alpha + 1, \quad \text{for all } \alpha < \kappa, \quad (1.2)$$

$$\|\mathbf{x} \vee \mathbf{y}\| = \|\mathbf{x}\| \vee \|\mathbf{y}\|, \quad \text{for all } \mathbf{x}, \mathbf{y} \in Q. \quad (1.3)$$

Now pick a partition $\kappa = \bigcup_{\alpha < \kappa} Z_\alpha$ of κ into sets Z_α such that $|Z_\alpha| = \kappa$ for all $\alpha < \kappa$. Define ideals I , I_0 , and I_1 of the Boolean algebra $B = P(\kappa)$ as follows:

$$\begin{aligned} I &= \{X \subseteq \kappa \mid X \text{ finite}\}, \\ I_0 &= \text{ideal of } B \text{ generated by } \{Z_\alpha \mid \alpha < \kappa\}, \\ I_1 &= \{X \subseteq \kappa \mid X \cap Z_\alpha \text{ is finite for every } \alpha < \kappa\}. \end{aligned}$$

It is obvious that $I = I_0 \cap I_1$, and that $\kappa \notin I_0 \cup I_1$. We define a map $\mu: B \rightarrow S$ by the following rule:

$$\mu(X) = \begin{cases} \bigvee_{\alpha \in X} \mathbf{q}_\alpha, & \text{if } X \text{ is finite,} \\ \mathbf{a}_\ell, & \text{if } X \in I_\ell \setminus I, \text{ for } \ell < 2, \\ \mathbf{a}_0 \vee \mathbf{a}_1, & \text{if } X \notin I_0 \cup I_1. \end{cases}$$

So μ is a $\{\vee, 0\}$ -homomorphism from B to S with $\mu(1) = \mathbf{a}_0 \vee \mathbf{a}_1$.

Now suppose that $\mu_0, \mu_1: B \rightarrow S$ satisfy (i)–(iii) above. For $\alpha < \kappa$, $\mu_1(Z_\alpha) \leq \mu(Z_\alpha) = \mathbf{a}_0$ (because $Z_\alpha \in I_0 \setminus I$), and $\mu_1(Z_\alpha) \leq \mu_1(\kappa) \leq \mathbf{a}_1$ (by the assumption (iii)), hence $\mu_1(Z_\alpha) \in Q$. Hence, since Z_α is a cofinal subset of κ , there exists $\xi_\alpha \in Z_\alpha$ such that $\alpha \vee \|\mu_1(Z_\alpha)\| \leq \xi_\alpha$. We put $Z = \{\xi_\alpha \mid \alpha < \kappa\}$. Observe that $Z \in I_1 \setminus I$, hence $\mu(Z) = \mathbf{a}_1$. So $\mu_0(Z) \leq \mu(Z) = \mathbf{a}_1$ on the one hand, and $\mu_0(Z) \leq \mu_0(\kappa) = \mathbf{a}_0$ on the other hand, thus $\mu_0(Z) \in Q$. Put $\beta = \|\mu_0(Z)\|$. Then

$$\begin{aligned} \xi_\beta + 1 &= \|\mathbf{q}_{\xi_\beta}\| && \text{(by (1.2))} \\ &= \|\mu(\{\xi_\beta\})\| && \text{(by the definition of } \mu) \\ &= \|\mu_0(\{\xi_\beta\})\| \vee \|\mu_1(\{\xi_\beta\})\| && \text{(by (i) and (1.3))} \\ &\leq \|\mu_0(Z)\| \vee \|\mu_1(Z_\beta)\| && \text{(by (ii))} \\ &= \beta \vee \|\mu_1(Z_\beta)\| \\ &\leq \xi_\beta, \end{aligned}$$

a contradiction. $\square$

In order to formulate Corollary 1.3, we recall the following definition, used in particular in [9]. It generalizes the classical definition of a weakly distributive homomorphism presented in [7].

Definition 1.2. Let S and T be join-semilattices, let $a \in S$. A join-homomorphism $\mu: S \rightarrow T$ is *weakly distributive at a* , if for all $b_0, b_1 \in T$ such that $\mu(a) = b_0 \vee b_1$, there are $a_0, a_1 \in S$ such that $a = a_0 \vee a_1$ and $\mu(a_\ell) \leq b_\ell$ for all $\ell < 2$.

Corollary 1.3. *Let S be a $\{\vee, 0\}$ -semilattice that is not a lattice. There exist a Boolean algebra B and a $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c B \rightarrow S$ such that there are no lattice L , no lattice homomorphism $f: B \rightarrow L$ and no $\{\vee, 0\}$ -homomorphism $\alpha: \text{Con}_c L \rightarrow S$ that satisfy the following properties:*

- (i) α is weakly distributive at $\Theta_L(f(0_B), f(1_B))$.
- (ii) $\varphi = \alpha \circ \text{Con}_c f$.

Proof. By assumption, there exist $\mathbf{a}_0, \mathbf{a}_1 \in S$ such that $Q = (\mathbf{a}_0] \cap (\mathbf{a}_1]$ has no largest element. We consider B, μ as in Lemma 1.1. Since the lattice B is Boolean, the rule $x \mapsto \Theta_B(0_B, x)$ defines an isomorphism $\pi: B \rightarrow \text{Con}_c B$. We put $\varphi = \mu \circ \pi^{-1}$.

So suppose that L, f , and α are as above. Observe that

$$\alpha\Theta_L(f(0_B), f(1_B)) = \alpha \circ (\text{Con}_c f)(\Theta_B(0_B, 1_B)) = \varphi\Theta_B(0_B, 1_B) = \mu(1_B) = \mathbf{a}_0 \vee \mathbf{a}_1,$$

thus, since α is weakly distributive at $\Theta_L(f(0_B), f(1_B))$, there are $\Psi_0, \Psi_1 \in \text{Con}_c L$ such that $\Psi_0 \vee \Psi_1 = \Theta_L(f(0_B), f(1_B))$ and $\alpha(\Psi_\ell) \leq \mathbf{a}_\ell$, for all $\ell < 2$. Thus there are a positive integer n and a decomposition

$$f(0_B) = t_0 \leq t_1 \leq \dots \leq t_{2n} = f(1_B) \quad (1.4)$$

in L such that the relations

$$\begin{aligned} t_{2i} &\equiv t_{2i+1} \pmod{\Psi_0}, \\ t_{2i+1} &\equiv t_{2i+2} \pmod{\Psi_1} \end{aligned}$$

hold for all $i < n$. For $x \in B$, we put

$$\begin{aligned} \mu_0(x) &= \bigvee_{i < n} \alpha\Theta_L(t_{2i} \wedge f(x), t_{2i+1} \wedge f(x)), \\ \mu_1(x) &= \bigvee_{i < n} \alpha\Theta_L(t_{2i+1} \wedge f(x), t_{2i+2} \wedge f(x)). \end{aligned}$$

We verify that conditions (i)–(iii) of Lemma 1.1 are satisfied, thus causing a contradiction.

Condition (i). For $x \in B$, we get

$$\begin{aligned} \mu_0(x) \vee \mu_1(x) &= \bigvee_{i < 2n} \alpha\Theta_L(t_i \wedge f(x), t_{i+1} \wedge f(x)) \\ &= \alpha\Theta_L(f(0_B) \wedge f(x), f(1_B) \wedge f(x)) && \text{(by (1.4))} \\ &= \alpha\Theta_L(f(0_B), f(x)) \\ &= \varphi(\Theta_B(0, x)) \\ &= \mu(x). \end{aligned}$$

Condition (ii). For $x \leq y$ and $i < n$, the relation

$$\Theta_L(t_{2i} \wedge f(x), t_{2i+1} \wedge f(x)) \subseteq \Theta_L(t_{2i} \wedge f(y), t_{2i+1} \wedge f(y))$$

holds (because $f(x) \leq f(y)$), thus $\mu_0(x) \leq \mu_0(y)$. So μ_0 is order-preserving. The proof that μ_1 is order-preserving is similar.

Condition (iii). For $i < n$, $\Theta_L(t_{2i}, t_{2i+1}) \subseteq \Psi_0$, thus $\alpha\Theta_L(t_{2i}, t_{2i+1}) \leq \alpha(\Psi_0) \leq \mathbf{a}_0$, whence $\mu_0(1) = \bigvee_{i < n} \alpha\Theta_L(t_{2i}, t_{2i+1}) \leq \mathbf{a}_0$. Similarly, $\mu_1(1) \leq \mathbf{a}_1$.

This contradicts, by Lemma 1.1, the existence of L, f , and α . $\square$

Theorem A. *Let S be a distributive $\{\vee, 0\}$ -semilattice. Then the following are equivalent:*

- (i) *For any lattice K and any $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c K \rightarrow S$, there are a lattice L , a lattice homomorphism $f: K \rightarrow L$, and an isomorphism $\alpha: \text{Con}_c L \rightarrow S$ such that $\varphi = \alpha \circ \text{Con}_c f$.*
- (ii) *S is a lattice.*

Proof. (ii) $\Rightarrow$ (i) follows from Theorem C in [10].

(i) $\Rightarrow$ (ii) is a particular case of Corollary 1.3. $\square$

With the terminology mentioned in the Introduction, this proves that 1-CLP holds at S iff S is a lattice, for any distributive $\{\vee, 0\}$ -semilattice S .

We also mention the following immediate consequence of Corollary 1.3, that solves (positively) the problem, stated by Dobbertin in [2], whether “strongly measurable semilattices are lattices”:

Corollary 1.4. *Let S be a distributive $\{\vee, 0\}$ -semilattice. Then the following are equivalent:*

- (i) *For any Boolean algebra B , any $\{\vee, 0\}$ -homomorphism $\mu: B \rightarrow S$, and any $\mathbf{a}_0, \mathbf{a}_1 \in S$ such that $\mu(1_B) = \mathbf{a}_0 \vee \mathbf{a}_1$, there are $\{\vee, 0\}$ -homomorphisms $\mu_0, \mu_1: B \rightarrow S$ such that $\mu = \mu_0 \vee \mu_1$ and $\mu_\ell(1_B) = \mathbf{a}_\ell$, for all $\ell < 2$.*
- (ii) *S is a lattice.*

Proof. (ii) $\Rightarrow$ (i) is proved in Corollary 10 of [1], see also [2].

(i) $\Rightarrow$ (ii) follows immediately from Corollary 1.3. $\square$

2. A COUNTEREXAMPLE OF SIZE $\aleph_1$

Throughout this section, we shall denote by D the $\{\vee, 0\}$ -semilattice defined as $D = \omega \cup \{\mathbf{a}_0, \mathbf{a}_1, \infty\}$, with ω a $\{\vee, 0\}$ -subsemilattice of D , $n < \mathbf{a}_\ell < \infty$ for all $\ell < 2$, and $\infty = \mathbf{a}_0 \vee \mathbf{a}_1$, see Figure 1.




FIGURE 1. The semilattice D

Now we shall construct a Boolean algebra B . By Cantor's Theorem, $\aleph_1 \leq 2^{\aleph_0}$, thus there exists a one-to-one map $f: \omega_1 \hookrightarrow \mathcal{P}(\omega)$ (where $\mathcal{P}(\omega)$ denotes the powerset of ω). We define a map $g: \omega_1 \times \omega_1 \rightarrow \omega$ by the rule

$$g(\xi, \eta) = \begin{cases} \text{least } n < \omega \text{ such that } f(\xi) \cap (n+1) \neq f(\eta) \cap (n+1), & \text{if } \xi \neq \eta, \\ 0, & \text{if } \xi = \eta. \end{cases}$$

Lemma 2.1. *Let $n < \omega$, let X be a subset of ω_1 . If $g(\xi, \eta) < n$ for all $\xi, \eta \in X$, then $|X| \leq 2^n$.*

Proof. Let p be the map from X to $\mathcal{P}(n)$ defined by the rule

$$p(\xi) = f(\xi) \cap n, \quad \text{for all } \xi \in X.$$

(We identify n with $\{0, 1, \dots, n-1\}$.) If $|X| > 2^n$, then there are $\xi, \eta \in X$ such that $\xi \neq \eta$ and $p(\xi) = p(\eta)$. Hence $g(\xi, \eta) \geq n$, by the definition of g , a contradiction. $\square$

Definition 2.2. We denote by B the Boolean algebra defined by generators $u_{0,\xi}$ and $u_{1,\xi}$, for $\xi < \omega_1$, and v_n , for $n < \omega$, and the following relations:

$$u_{0,\xi} \wedge u_{1,\eta} \leq v_{g(\xi,\eta)}, \quad \text{for all } \xi, \eta < \omega_1. \quad (2.1)$$

Furthermore, we put $w_n = \bigvee_{k \leq n} v_k$, for all $n < \omega$.

Lemma 2.3. $u_{0,\xi} \wedge u_{1,\eta} \leq w_n$ iff $g(\xi, \eta) \leq n$, for all $\xi, \eta < \omega_1$ and all $n < \omega$.

Proof. If $g(\xi, \eta) \leq n$, then $u_{0,\xi} \wedge u_{1,\eta} \leq w_n$ by (2.1).

Conversely, suppose that $u_{0,\xi} \wedge u_{1,\eta} \leq w_n$. We define elements $u_{0,\xi'}^*$, $u_{1,\eta'}^*$, and v_k^* of the two-element Boolean algebra $\mathbf{2}$, for $\xi', \eta' < \omega_1$ and $k < \omega$, as follows:

$$u_{0,\xi}^* = u_{1,\eta}^* = 1; \quad (2.2)$$

$$u_{0,\xi'}^* = 0, \quad \text{for all } \xi' < \omega_1 \text{ such that } \xi' \neq \xi; \quad (2.3)$$

$$u_{1,\eta'}^* = 0, \quad \text{for all } \eta' < \omega_1 \text{ such that } \eta' \neq \eta; \quad (2.4)$$

$$v_{g(\xi,\eta)}^* = 1; \quad (2.5)$$

$$v_k^* = 0, \quad \text{for all } k < \omega \text{ such that } k \neq g(\xi, \eta). \quad (2.6)$$

Let $\xi', \eta' < \omega_1$. If $\xi' = \xi$ and $\eta' = \eta$, then $u_{0,\xi'}^* \wedge u_{1,\eta'}^* = 1 = v_{g(\xi,\eta)}^*$. Otherwise, $u_{0,\xi'}^* \wedge u_{1,\eta'}^* = 0 \leq v_{g(\xi',\eta')}^*$. So the elements $u_{0,\xi'}^*$, $u_{1,\eta'}^*$, and v_k^* , for $\xi', \eta' < \omega_1$ and $k < \omega$, verify the inequalities (2.1). Therefore, there exists a homomorphism of Boolean algebras $\varphi: B \rightarrow \mathbf{2}$ such that

$$\varphi(u_{\ell,\xi'}) = u_{\ell,\xi'}^*, \quad \text{for all } \xi' < \omega_1 \text{ and } \ell < 2,$$

$$\varphi(v_k) = v_k^*, \quad \text{for all } k < \omega.$$

In particular, by assumption, $u_{0,\xi}^* \wedge u_{1,\eta}^* \leq \bigvee_{k \leq n} v_k^*$, that is, $\bigvee_{k \leq n} v_k^* = 1$. Therefore, by (2.6), $g(\xi, \eta) \leq n$. $\square$

Theorem B. *There exist a Boolean algebra B of size $\aleph_1$ and a $\{\vee, 0\}$ -homomorphism $\mu: B \rightarrow D$ such that the following holds:*

- (a) $\mu(1_B) = \infty$;
- (b) *there are no maps $\mu_0, \mu_1: B \rightarrow D$ that satisfy the following properties:*
 - (i) $\mu(x) = \mu_0(x) \vee \mu_1(x)$, for all $x \in B$,
 - (ii) μ_0 and μ_1 are order-preserving,
 - (iii) $\mu_\ell(1) \leq \mathbf{a}_\ell$, for all $\ell < 2$.

Proof. Let B be the Boolean algebra constructed in Definition 2.2. It is clear that $|B| = \aleph_1$. We define ideals I_0 , I_1 , and I of B , as follows:

$$I_\ell = \text{ideal of } B \text{ generated by } \{u_{\ell,\xi} \mid \xi < \omega_1\} \cup \{v_k \mid k < \omega\}, \text{ for all } \ell < 2,$$

$$I = \text{ideal of } B \text{ generated by } \{v_k \mid k < \omega\}.$$

It follows from (2.1) that $I_0 \cap I_1 = I$. Therefore, we can define a $\{\vee, 0\}$ -homomorphism $\mu: B \rightarrow D$ by the rule

$$\mu(x) = \begin{cases} \text{least } n < \omega \text{ such that } x \leq w_n, & \text{if } x \in I_0 \cap I_1, \\ \mathbf{a}_\ell, & \text{if } x \in I_\ell \setminus I_{1-\ell}, \text{ for } \ell < 2, \\ \infty, & \text{if } x \notin I_0 \cup I_1, \end{cases}$$

for all $x \in B$.

Now let $\mu_0, \mu_1: B \rightarrow D$ satisfying (i)–(iii) above. We put

$$\begin{aligned} X_n &= \{ \xi < \omega_1 \mid \mu_1(u_{0,\xi}) \leq n \}, \\ Y_n &= \{ \eta < \omega_1 \mid \mu_0(u_{1,\eta}) \leq n \}, \end{aligned}$$

for all $n < \omega$.

Claim 1.

- (a) *The sequences $\langle X_n \mid n < \omega \rangle$ and $\langle Y_n \mid n < \omega \rangle$ are increasing.*
- (b) $\omega_1 = \bigcup_{n < \omega} X_n = \bigcup_{n < \omega} Y_n$.

Proof of Claim. (i) is trivial.

(ii) Let $\xi < \omega_1$. Then

$$\begin{aligned} \mu_1(u_{0,\xi}) &\leq \mu(u_{0,\xi}) && \text{(by assumption (i))} \\ &\leq \mathbf{a}_0 && \text{(by the definition of } \mu), \end{aligned}$$

while also $\mu_1(u_{0,\xi}) \leq \mathbf{a}_1$ by assumptions (ii) and (iii). Therefore, $\mu_1(u_{0,\xi}) \leq n$ for some $n < \omega$. This proves that $\omega_1 = \bigcup_{n < \omega} X_n$. The proof that $\omega_1 = \bigcup_{n < \omega} Y_n$ is similar. $\square$ Claim 1.

Now we put $Z_n = X_n \cap Y_n$, for all $n < \omega$. It follows from Claim 1 that $\omega_1 = \bigcup_{n < \omega} Z_n$. In particular, one of the Z_n should be infinite (and even uncountable). We fix such an n . For all $\xi, \eta \in Z_n$, $\mu_1(u_{0,\xi}) \leq n$ and $\mu_0(u_{1,\eta}) \leq n$, thus, by assumptions (i) and (ii), $\mu(u_{0,\xi} \wedge u_{1,\eta}) \leq n$, that is, $u_{0,\xi} \wedge u_{1,\eta} \leq w_n$. Thus, by Lemma 2.3, $g(\xi, \eta) \leq n$. Hence, by Lemma 2.1, Z_n is finite, a contradiction. $\square$

Corollary 2.4. *There exist a Boolean algebra B of size $\aleph_1$ and a $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c B \rightarrow D$ such that there are no lattice L , no lattice homomorphism $f: B \rightarrow L$ and no $\{\vee, 0\}$ -homomorphism $\alpha: \text{Con}_c L \rightarrow D$ that satisfy the following properties:*

- (i) α is weakly distributive at $\Theta_L(f(0_B), f(1_B))$.
- (ii) $\varphi = \alpha \circ \text{Con}_c f$.

Proof. As in the proof of Corollary 1.3. $\square$

3. OPEN PROBLEMS

The main result of Theorem A states that the possibility, for a given distributive $\{\vee, 0\}$ -semilattice S , to lift every $\{\vee, 0\}$ -homomorphism $\text{Con}_c K \rightarrow S$ for any lattice K is equivalent to S being a lattice. The maps considered in the proof of this result are not one-to-one. This leaves open the following question:

Problem 1. Let S be a distributive $\{\vee, 0\}$ -semilattice. When is it possible to lift every *one-to-one* $\{\vee, 0\}$ -homomorphism $\varphi: \text{Con}_c K \hookrightarrow S$, for any lattice K ?

By Theorem C of [10], the condition that S be a lattice is sufficient. Is this condition also necessary?

Problem 2. Let K be a lattice, let S be a distributive $\{\vee, 0\}$ -semilattice, let $\varphi: \text{Con}_c K \rightarrow S$ be a *distributive* $\{\vee, 0\}$ -homomorphism. Can φ be lifted?

Recall (see [7]) that for $\{\vee, 0\}$ -semilattices S and T , a homomorphism $\varphi: S \rightarrow T$ is *distributive*, if φ is *surjective* and $\ker \varphi$ is a directed union of the form $\bigcup_{i \in I} \ker s_i$, where s_i is a closure operator on S for all i . The result of Corollary 1.3 is of no help for solving Problem 2, because the contradiction follows there from the failure of α to be (weakly) distributive.

Problem 3. Let K be a countable lattice, let S be a countable distributive $\{\vee, 0\}$ -semilattice. Can every $\{\vee, 0\}$ -homomorphism from $\text{Con}_c K$ to S be lifted?

For countable S , not every $\{\vee, 0\}$ -homomorphism from $\text{Con}_c K$ to S can be lifted as a rule, even for K of size $\aleph_1$ (this follows from Corollary 2.4). However, the problem is still open for countable K .

Our last problem is more oriented to axiomatic set theory. It originates in the observation that the construction of the Boolean algebra of the proof of Theorem A does not rely on the Axiom of Choice (but it has size the continuum), while the construction of the Boolean algebra of the proof of Theorem B does not rely on the Continuum Hypothesis (but it relies on the Axiom of Choice, in the form of the existence of a one-to-one map from ω_1 into $\mathcal{P}(\omega)$).

Problem 4. Can one prove Theorem B by using neither the Axiom of Choice nor the Continuum Hypothesis?

REFERENCES

- [1] H. Dobbertin, *Vaught measures and their applications to lattice theory*, J. Pure Appl. Algebra **43** (1986), 27–51.
- [2] ———, “Boolean Representations of Refinement Monoids and Their Applications”, manuscript.
- [3] G. Grätzer, “General Lattice Theory. Second Edition”, Birkhäuser Verlag, Basel, 1998. xix+663 pp.
- [4] G. Grätzer, H. Lakser, and E.T. Schmidt, *Representing isotone maps as maps of congruences. I. Abstract maps*, Acta Math. Sci. Hungar. **75** (1997), 105–135.
- [5] ———, *Representing isotone maps as maps of congruences. II. Concrete maps*, preprint.
- [6] G. Grätzer and E.T. Schmidt, *Congruence Lattices*, Appendix C in [3], 519–530.
- [7] E.T. Schmidt, *Zur Charakterisierung der Kongruenzverbände der Verbände*, Mat. Časopis Sloven. Akad. Vied **18** (1968), 3–20.
- [8] J. Tůma and F. Wehrung, *Simultaneous representations of semilattices by lattices with permutable congruences*, Internat. J. Algebra Comput., to appear.
- [9] F. Wehrung, *A uniform refinement property for congruence lattices*, Proc. Amer. Math. Soc. **127**, no. 2 (1999), 363–370.
- [10] ———, *Forcing extensions of partial lattices*, manuscript.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, SOKOLOVSKÁ 83, CHARLES UNIVERSITY, 186 00 PRAHA 8, CZECH REPUBLIC
E-mail address: tuma@karlin.mff.cuni.cz

CNRS, FRE 2271, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE CAEN, 14032 CAEN CEDEX, FRANCE
E-mail address: wehrung@math.unicaen.fr
URL: <http://www.math.unicaen.fr/~wehrung>